

Edge effect on electronic properties in nanoscale graphene systems

Wakabayashi K.^{1,2}

¹ WPI-MANA, NIMS, 305-0044, Tsukuba, Japan

² PRESTO, JST, Japan

e-mail: WAKABAYASHI.Katsunori@nims.go.jp

The discovery of graphene and successive fabrication of graphene devices [1,2] have triggered intensive and diverse research on carbon related systems. The honeycomb crystal structure of single layer graphene consists of two nonequivalent sublattices and results in a unique band structure for the itinerant p-electrons near the Fermi energy which behave as massless Dirac fermion. In graphene, the presence of edges can have strong implications for the spectrum of the p-electrons. In graphene nanoribbons with zigzag edges, localized states appear at the edge with energies close to the Fermi level.[3] In contrast, edge states are absent for ribbons with armchair edges. Recent experiments have succeeded to synthesize graphene nanoribbons using lithography techniques[4], chemical techniques.[5,6]

In my talk, we focus on edge and geometry effects of the electronic properties of graphene nanoribbons. The electronic states of graphene nanoribbons crucially depend on the edge orientation and boundary condition.[3,7] (1) In zigzag nanoribbons, for disorder without inter-valley scattering a single perfectly conducting channel emerges associated with such a chiral mode due to edge states, i.e. the absence of the localization.[8-10] (2) In armchair nanoribbons, the single-channel transport subjected to long-ranged impurities is nearly perfectly conducting, where the backward scattering matrix elements in the lowest order vanish as a manifestation of the existence of Berry phase.[11-12] (3) Nano-graphene junctions are shown to have the zero-conductance anti-resonances associated with the edge states. The relation between the condition of the resonances and geometry is discussed.[13] (4) Finally, we will discuss the effect of edge chemical modification on magnetic properties of nanographene systems[14], and the hole doping effect on zigzag edge magnetic states.[15]

- [1] K.S. Novoselov, A.K. Geim, S.V. Morozov, et.al., *Nature* **438**, 197 (2005).
- [2] Y. Zhang, Y.-W.Tan, H.L.Stormer, and P.Kim, *Nature* **438**, 201 (2005).
- [3] M. Fujita, K. Wakabayashi, et.al, *J. Phys. Soc. Jpn.* **65**, 1920 (1996).
- [4] M.Y. Han, B. Oezylmaz, Y. Zhang, and P. Kim, *Phys. Rev. Lett.* **98**, 206805 (2007).
- [5] X. Li, X. Wang, L. Zhang, S. Lee, H. Dai, *Science* **31**, (2008) 122.
- [6] J. Cai, P. Ruffieux, R. Jaafar, M. Bieri, et.al., *Nature* **466**, 470 (2010).
- [7] K. Wakabayashi, et.al., *Sci. Technol. Adv. Mat.* **11**, 054504 (2010).
- [8] K. Wakabayashi, et.al., *Phys. Rev. Lett.* **99**, 036601 (2007).
- [9] K. Wakabayashi, et.al., *CARBON* **47** (2009) 124
- [10] K. Wakabayashi, *New J. Phys.* **11** (2009) 095016.
- [11] M. Yamamoto, Y. Takane, and K. Wakabayashi, *Phys. Rev.* **B79** (2009) 125421.
- [12] K. Sasaki, K. Wakabayashi, T. Enoki, *New J. Phys.* **12** (2010) 083023.
- [13] M. Yamamoto and K. Wakabayashi, *Appl. Phys. Lett.* **95** (2009) 082109.
- [14] K. Wakabayashi, S. Okada et.al., *J. Phys. Soc. Jpn.* **79** (2010) 034706
- [15] S. Dutta and K. Wakabayashi, in preparation.